WHY IT'S IMPORTANT TO GET VACCINATED FOR HEPATITIS B

The CDC recommends hepatitis B vaccination for all adults aged 19-59*

WHAT YOU NEED TO KNOW

Hepatitis B is a vaccine-preventable liver infection caused by the hepatitis B virus (HBV)



HBV is a leading cause of cirrhosis and liver cancer globally



There is no cure for hepatitis B. Vaccination is the best way to prevent hepatitis B



HBV is up to 100 times more infectious than HIV



The threat can linger— HBV can live on surfaces for at least 7 days



~296 million people are living with hepatitis B[†]

GETTING PROTECTED

HEPLISAV-B is the only adult hepatitis B vaccine that protects you with just 2 doses in 1 month



* The CDC also recommends hepatitis B vaccination for adults aged >60 with risk factors. Adults aged >60 without risk factors may receive hepatitis B vaccination.

+ Estimate based on World Health Organization data from 2019.

CDC, Centers for Disease Control and Prevention.

INDICATION

HEPLISAV-B is indicated for the prevention of infection caused by the hepatitis B virus in adults 18 years of age and older.

SELECT IMPORTANT SAFETY INFORMATION

If you have a history of severe allergic reaction after a previous dose of any hepatitis B vaccine, or to any ingredient of HEPLISAV-B, including yeast, do not take HEPLISAV-B.

Please see additional Important Safety Information on next page and <u>click here</u> for full Prescribing Information.





REMEMBER TO SCHEDULE YOUR SECOND SHOT FOR A MONTH LATER. It's important that you receive both shots to be fully protected.

TEXT HEPB TO 1-844-402-4372 and we'll send you a reminder when you're due for your final dose of HEPLISAV-B.

HEPLISAV-B has the same insurance coverage as all other CDC-recommended adult vaccines.

Learn more at **HEPLISAVB.com**

INDICATION

HEPLISAV-B is indicated for the prevention of infection caused by the hepatitis B virus in adults 18 years of age and older.

IMPORTANT SAFETY INFORMATION

If you have a history of severe allergic reaction after a previous dose of any hepatitis B vaccine, or to any ingredient of HEPLISAV-B, including yeast, do not take HEPLISAV-B.

HEPLISAV-B must be given by a medical professional, who will monitor you afterwards to check for allergic reaction.

If you are immunocompromised, or receiving immunosuppressant therapy, you may have less of an immune response to HEPLISAV-B.

Some people have hepatitis B infection without being aware of it or showing any symptoms. If you already have hepatitis B present in your body, HEPLISAV-B may not prevent hepatitis B infection.

The most common side effects reported by patients within 7 days of vaccination with HEPLISAV-B were pain at the site of injection (23%-39%), tiredness (11%-17%), and headache (8%-17%).

Please <u>click here</u> for full Prescribing Information.



